

Common Bacterial Blight



Photograph provided by H. F. Schwartz
AglImage - Colorado State University

Common bacterial blight (CBB) caused by *Xanthomonas campestris* pv. *phaseoli* (Smith) Dowson is a serious seed-borne disease of beans in both temperate and tropical production zones. Management of common blight includes the use of disease-free seed, crop rotation and resistant varieties (Saettler, 1989).

Zapata et al. (1985) prepared inoculum for greenhouse screening using cell suspensions of *Xanthomonas campestris* cultures grown at 48 h at 26° C. All concentrations were adjusted to 10^9 cfu/ml using standard turbidometric and dilution plating techniques. Inoculum for field inoculation was prepared by collecting leaves from plants at the R6-R7 stage of development with typical common bacterial blight symptoms. The infected leaves were macerated in a high-speed blender with 10 ml of 0.01 M phosphate buffer (pH 7) per 1 g of infected tissue. In the greenhouse, trifoliolate leaves of plants at initiation of flowering (R1 stage of development) were inoculated by placing the bean leaf over a plastic sponge saturated with inoculum and perforating it with a multiple-neededled florist frog (2 cm square metal base supporting rows of needles 3 mm apart and 12 mm in length). After inoculation, high humidity was maintained by mist irrigation every few minutes. In the field, a pneumatic sandblasting pistol connected by an air hose to an air compressor was used to inoculate bean plants at the R1 stage of development. The bacterial suspension from the field-infected leaves and 28.4 g of fine silica sand was placed in the reservoir of the pistol. The mixture was sprayed twice over one-half of each row at 345 kPa (50 psi) with the pistol held 30-60 cm from the plant. This method differs from the technique reported by Webster et al. (1983b) who sprayed a suspension of 5×10^7 cfu/ml onto plants using a sprayer nozzle at 3 kg/cm² at weekly intervals beginning 3 weeks after planting.

Gilbertson et al. (1988) prepared dry leaf inoculum from diseased bean leaves collected 14 days after greenhouse plants had been razor-blade-inoculated. Diseased leaves were dried at 28-30° C for 24-48 h, ground thoroughly in a Waring blender, sifted through a 32 mesh sieve (< 500 µm) and stored in plastic bags at 4 or -20° C. Pathogenic strains of *Xanthomonas campestris* pv. *phaseoli* were recovered from dry-leaf inoculum stored up to 6 years. Dry leaf inoculum can also be prepared using leaves collected from field from plants expressing common bacterial blight symptoms (Saettler, 1989). To prepare the suspension, 4 g of dry leaf inoculum was mixed with 1.9 L of water plus three drops of sticker

Zapata et al. (1985) prepared inoculum for the evaluation of pods using cell suspensions of *Xanthomonas campestris* cultures grown at 48 h at 26° C. All concentrations were adjusted to 10⁹ cfu/ml using standard turbidometric and dilution plating techniques. Pods in mid-pod-fill were harvested from plants, disinfected in a dilute sodium hypochlorite solution (0.02% Cl) and washed three times in tap water. The pods were placed on moistened paper towels in trays. Two needles mounted 2 mm apart were used to make 4-5 mm long scratches through the epidermis of the pods. A 0.01 mm drop of inoculum was deposited over the distal scratch marks. The pods were incubated in an environmental chamber at 27° C with fluorescent lighting (3,220 lux) for 12 h. Readings were taken daily. A 1-5 scale was used to evaluate the pods, where 1 = no water soaking, 2 = 1mm wide water soaked area, 3 = 2-3 mm wide water soaked area, 4 = water soaked region coalescing over scratches and 5 = water soaked area reached the pod suture. Ariyaratne et al (1998) observed similar symptoms for detached and attached pods inoculated with *Xcp* and suggested using the former to reduce environmental variance.

Varela et al. (1996) evaluated two common bean populations for CBB resistance in the field in the F₃ and F₄ generations and in the greenhouse in the F₅ generation. A randomized complete block design with three replications was used. Experimental units were 1 m row lengths in the field and single pots in the greenhouse. The lines were inoculated in the field at the R6 stage of development with a backpack sprayer using a bacterial concentration of 3 x 10⁷ cfu/ml (Schuster and Coyne, 1981). In the greenhouse, the lines were inoculated at 14 days after planting using the multiple needle technique. The lines were evaluated 14 days after inoculation for CBB reaction using the CIAT 1-9 scale and percentage of leaf area affected. Selection in the F₄ and F₅ generations was more effective because of higher heritabilities. CBB evaluations using the CIAT 1-9 scale were better than estimates of percentage of leaf area infected due to higher heritabilities and greater precision to detect differences among lines.

Singh and Muñoz (1999) developed dry beans at CIAT in Colombia with high levels of resistance to common bacterial blight by field screening advanced generation (≥ F₅) lines in unreplicated single row plots. Susceptible and resistant checks having known CBB reactions were planted throughout the nursery.

Beginning about 3 weeks after planting, the rows were inoculated two to four times at 7-10 day intervals. The leaves in the rows were inoculated with a backpack sprayer using a bacterial concentration between 10^7 and 10^8 cfu/mL. The most CBB resistant lines were subsequently evaluated in replicated trials. Three leaflets of the first or second trifoliate leaf of the resistant lines were inoculated with twin surgical blades about 3 weeks after planting. The lines were also inoculated 3-5 times, at weekly intervals, by spraying the canopy. At mid-pod-fill the pods were inoculated with multiple needles using a florist frog. The trifoliate leaves were evaluated 7 to 15 days after inoculation and the pods were evaluated about 1 week after inoculation.

Singh and Muñoz (1999) noted several problems related to breeding for CBB resistance. Single plant selection under high disease pressure is recommended to maintain high levels of CBB resistance. Differential CBB reaction of leaves and pods requires that both plant organs must be screened for resistance. Because resistance to specific strains of *Xcp* has been observed, lines should be inoculated with strains from the target region. Finally, plants should be evaluated for CBB reaction at the same stage of development to avoid selecting plants having later maturity.

Aggour et al. (1988a) evaluated pod wall and pedicel inoculation techniques to screen beans for resistance to common bacterial blight. The edge of a razor blade that had been immersed in a 48-hour culture of *Xcp* was used to inoculate bean pods (R7-R8). The pods were evaluated 15 days after inoculation using a 1-9 scale, where 1 = no visible lesion, 2 = < 0.5 - < 1.0 mm, 3 = 1.0 mm, 4 = 2 mm, 5 = 3 mm, 6 = 4 mm, 7 = 5 mm, 8 = 6 mm and 9 = \geq 7 mm sized lesions. The method was successful in detecting a differential pod reaction on bean lines. The pedicels of flower buds (24 to 36 hours before anthesis) and small pods (2-3 days old) were injected with about 0.01 ml of *Xcp* inoculum (10^8 cfu in a PO_4 buffer, pH 7.1) using a syringe with a 30 gage x 1.77 cm needle. When the pedicel was too thin, a surface scratch was made on the pedicel and a drop of inoculum was placed over the scratch. The technique was found to be useful to select for pedicel resistance of *Xcp*.

Aggour et al. (1988b) described different seed inoculation methods to select beans for resistance to common bacterial blight. A concentration of 10^6 cfu in a PO_4 buffer (pH 7.1) was used for all inoculation methods. Soaking dry seed in a bacterial cell suspension for 6 hours to inoculate the embryo produced the greatest number of seedlings with visible symptoms. There was good agreement between results of the seed inoculation methods and leaf inoculation of seedlings using the multi-needle method.

Resistant lines derived from the cross Tio Canela 75/VAX 6 were selected in Honduras for resistance to virulent isolates of *Xcp* using the multiple needle inoculation technique and marker-assisted selection (MAS) for the presence of the SCAR markers SAP6 and SU91. (Beaver and Rosas, unpublished).

Miklas et al. (2005) and Mutlu et al. (2005a and b) used field screening and multiple needle inoculations in the greenhouse, combined with MAS for the SCAR markers SAP6 and SU91, to develop CBB resistant dark red kidney (USDK-CBB-15) and pinto (ABCP-8) germplasm lines.

Park and Dhanvantari (1987) prepared inoculum of four local strains, two of *X. campestris* pv *phaseoli* var. *fuscans* no. 12 and no. 118 and two of non-*fuscans* no. 18 and no. 98 that were grown on yeast salts-agar (Dye 1968) for 48 h at 25° C. A sterile distilled water suspension of bacterial growth was made, adjusting the concentration to 15 Klett units against a red filter in a Klett-Summerson colorimeter. Suspensions of the four strains were mixed in equal volume and 7 L were added to 190 L of water in a spray tank. The inoculum, containing 10^6 colony-forming units mL^{-1} , was applied at about 1333 kPa to ensure visible water-soaking injury using a spray boom. Bean plants were inoculated at about 3 wk old in a field screening nursery. Assessment for blight was made twice, 3 and 6 wk after inoculation. Leaf blight was rated visually as: 1 = no visible symptoms, 2 = small flecks or lesions on less than 5 % of leaf surface, 3 = small lesions covering 5-20 % of leaf surface, 4 = lesions of various sizes covering 20-40% of leaf surface with some chlorosis, 5 = lesions coalescing to cover 40-60 % of leaf surface and surrounded by orange-yellow chlorosis, 6 = large necrotic lesions with spreading chlorosis covering 60-80 % of leaf surface, and 7 = lesions and chlorosis spreading, leaf wilting and defoliation. Leaf blight was scored on 10 single plants per row. Pod blight was rated as: 1 = no visible symptoms, 2 = a few small lesions, 3 = lesions of moderate size, 4 = large lesions, 5 = large coalescing lesions. Percent pod infection was recorded as number of infected pods out of total number of pods from plants per row. We also inoculated bean plants by the multiple needle method (Andrus, 1948) under controlled environmental condition in growth rooms/greenhouses.

Park and Dhanvantari (1993) developed common bean germplasm line HR45 (Reg. no. GP-114, PI 570661) at Agriculture and Agri-Food Canada Greenhouse & Processing Crops Research Centre, Harrow, ON, and released in 1993. It was selected for small white navy bean type with improved resistance to common bacterial blight incited by *Xanthomonas campestris* pv. *phaseoli* (Smith) Dye and for upright plant type. HR45 was derived from the cross of HR13-621*2//XAN159/HR13-621, of which the original single cross between XAN159 and HR13-621 was made in 1986.

Michaels et al. (2005) developed CBB resistant navy bean cv. OAC Rex, tested as OAC 95-4, which was derived from the cross HR20-728/MBE 7 made in 1988. MBE 7 was a selection from the cross ICA Pijao/PI 440795//Ex Rico 23 and was used to provide resistance to common bacterial blight, transferred through an interspecific cross, and subsequent embryo rescue, from the *Phaseolus acutifolius* A. Gray line PI 440795 (Parker, 1985). F_1 plants were grown in the

field in 1988 at the University of Guelph Elora Research Station (ERS), Elora, Ontario. F₆ plants were screened in an indoor growth room in the winter of 1991 for CBB resistance using multiple-pin techniques (Andrus, 1948). Resistant selections were grown in the field at ERS in 1991 and 1992 as F₆ and F₇ generations, respectively and assessed for CBB resistance using the dry leaf inoculum method (Gilbertson et al., 1988). OAC 95-4 was tested in Ontario coop cultivar registration and performance trials during 1995-1999. OAC Rex was registered (registration no. 5491) on May 24, 2002 by Variety Registration Office, Canadian Food Inspection Agency, AAFC.



Photograph provided by H. F. Schwartz
AgImage - Colorado State University

Table 1. Sources of resistance to common bacterial blight in different seed classes.

Name or number	Seed color / type	QTL marker & Resistance sources	Reference
VAX 5 ICB-3 USBK-CBB-5 XAN 112	9 / Black	SAP6, SU91 SAP6, P. coccineus SAP6 SAP6, PI207262	Singh et al. (1999) Miklas et al. (1999) Miklas et al. (2001a) CIAT
ICB-8 USNA-CBB-1 to 4 Wilk-2 CBB-Teebus HR-45 OAC Rex	1 / White	SAP6, P. coccineus SAP6 SAP6, SU91, & BC420 SU91, BC420 BC420 BC73	Miklas et al. (1999) Miklas et al. (2001a) Cornell Fourie and Herselman (2002) Park & Dhanvantari (1993) Michaels et al. (2005?)
USPT-CBB-1 to 3 Chase ABCP-8	2M / Pinto	SAP6, P. coccineus SAP6 SAP6, SU91	Miklas et al. (2001b) Coyne et al. (1994) Mutlu et al (2005)
USGN-CBB-4 Wei hing BelNeb-RR-1	1 / Great Northern	SAP6 SAP6	Miklas et al. (2001b) Coyne et al. (2000) Stavely et al. (1989)
VAX 6	7 / Purple	SAP6, SU91	Singh and Muñoz (1999)
VAX 3 ICB-6 XAN 309	6 / Small red	SAP6, SU91 SAP6, P. coccineus SAP6, SU91	Singh et al. (2001) Miklas et al. (1999) CIAT
	5 / Pink		
USCR-CBB-12 & 13	2R / Cranberry	SAP6	Miklas et al. (2001c)
Pomjor 17	6M / Red mottled		Beaver et al. (1992)
PR9443-4 USLK-CBB-9 & 10 USDK-CBB-11 Montcalm USDK-CBB-15	5K / Light red kidney 6K / Dark red kidney	 SAP6 SAP6, SU91	Beaver et al. (1999c) Miklas et al. (2001c) MSU (Adams, Saettler) Miklas et al. (2005)
	1 / Snap		
W-BB -11,-20-1, - 35, -52, -11-56	Miscellaneous	SAP6	Zapata et al. 2004

Molecular marker UBC420 for HR45 (Yu et al., 2000 and 2004)

Molecular marker BC73 for OAC Rex (Bai et al., 1997; Miklas, et al. 2000)

References

- Aggour, A.R., D.P. Coyne and A. Vidaver. 1988a. Inoculation techniques to evaluate reaction of beans (*Phaseolus vulgaris*) to *Xanthomonas campestris* pv. *phaseoli*. Ann. Rep. Bean Improv. Coop. 31:134-135.
- Aggour, A.R., D.P. Coyne and A.K. Vidaver. 1988b. Potential of seed inoculation to select for resistance to common blight in beans (*Phaseolus vulgaris* L.). Ann. Rep. Bean Improv. Coop. 31:136-137.
- Bai, Yonghe, T. E. Michaels and K. P. Pauls. 1997. Identification of RAPD markers linked to common bacterial blight resistant genes in *Phaseolus vulgaris* L. Genome 40: 544-551.
- Beaver, J.S., M. Zapata and P.N. Miklas. 1999c. Registration of PR9443-4, bean germplasm resistant to bean golden mosaic, common bacterial blight and rust. Crop Sci. 39:1262.
- Beaver, J.S., J.R. Steadman and D.P. Coyne. 1992. Field reaction of landrace components of red mottled beans to common bacterial blight. HortScience 27:50-51.
- Coyne, D.P., D.S. Nuland, D.S., D.T. Lindgren, and J.R. Steadman. 1994. >Chase= pinto dry bean. HortScience 29:44-45.
- Coyne DP, Nuland DS, Lindgren DT, Steadman JR, Smith DW, Gonzales J, Schild J, Reiser J, Sutton L, Carlson C, Stavely JR, and Miklas PN. 2000. Weihing great northern disease resistant dry bean. HortScience 35:310-312.
- Fourie, D. and L. Herselman. 2002. Breeding for common blight resistance in dry beans in South Africa. Annu. Rpt. Bean Improv. Coop. 45: 50-51.
- Gilbertson, R.L., R.E. Rand, E. Carlson and D.J. Hagedorn. 1988. The use of dry leaf inoculum for the establishment of common bacterial blight of beans. Plant Disease 72:385-389.
- Michaels, T.E., T. H. Smith and A. D. Beatie. 2005. OAC Rex common bean. Can. J. Plant Sci. (in press).
- Miklas, P.N., M. Zapata, J.S. Beaver and K.F. Grafton. 1999. Registration of four dry bean germplasms resistant to common bacterial blight: ICB-3, ICB-6, ICB-8 and ICB-10. Crop Sci. 39:594.
- Miklas, P.N., J.R. Smith, A.N. Hang, K.F. Grafton and J.D. Kelly. 2001a. Release of navy and black bean germplasm lines USNA-CBB-1 to 4 and USBK-CBB-5 with resistance to common bacterial blight. Ann. Rep. Bean Improv. Coop. 44:181-182.
- Miklas, P.N., J.R. Smith, K.F. Grafton, D.P. Coyne and M.A. Brick. 2001b. Release of pinto and great northern bean germplasm lines USPT-CBB-1 to 3 and USGN-CBB-4 with erectness and resistance to common bacterial blight, rust and mosaic. Ann. Rep. Bean Improv. Coop. 44:182-183.

Miklas, P.N., J.R. Smith and J.D. Kelly. 2001c. Release of kidney and cranberry dry bean germplasm lines USLK-CBB-9 & 10, USDK-CBB-11 and USCR-CBB-12 & 13 with resistance to common bacterial blight and anthracnose. Ann. Rep. Bean Improv. Coop. 44:186-188.

Miklas PN, J.R. Smith, and S.P. Singh. 2005. Release of common bacterial blight resistant dark red kidney bean germplasm line USDK-CBB-15. Ann Rep Bean Improv Coop. 48: 192-193.

Mutlu, N., P. Miklas, J. Reiser, and D. Coyne. 2005a. Backcross breeding for improved resistance to common bacterial blight in pinto bean (*Phaseolus vulgaris* L.). Plant Breeding 124, 282-287.

Mutlu, N., P.N. Miklas, J.R. Steadman, A.V. Vidaver, D. Lindgren, J. Reiser, and M.A. Pastor-Corrales. 2005b. Registration of pinto bean germplasm line ABCP-8 with resistance to common bacterial blight. Crop Sci. 45:806-807.

Park, S.J. and B. N. Dhanvantari. 1989. Transfar of common bacterial blight (*Xanthomona campestris* pv *phaseoli*) resistance from *Phaseolus coccineus* Lam. to *P. vulgaris* L. through interspecific hybridization. Can. J. Plant Sci. 67: 685-695.

Park, S.J. and B. N. Dhanvantari. 1994. Registration of common bean blight-resistant germplasm, HR45. Crop Sci. 34: 548.

Saettler, A.W. 1989. Common bacterial blight. p. 261-284. In H.F. Schwartz and M.A. Pastor-Corrales (eds.). Bean Production Problems in the Tropics. Centro Internacional de Agricultura Tropical (CIAT). Cali, Colombia.

Schuster, M.L. and D.P. Coyne. 1981. Biology, epidemiology, genetics and breeding for resistance to bacterial pathogens of *Phaseolus vulgaris* L. Hort. Rev. 3:28-58.

Singh, S.P., C.G. Muñoz and H. Terán. 2001. Registration of common bacterial blight resistant dry bean germplasm VAX 1, VAX 3 and XAX 4. Crop Sci. 41:275-276.

Singh, S.P. and C.G. Muñoz. 1999. Resistance to common bacterial blight among *Phaseolus* species and common bean improvement. Crop Sci. 39:80-89.

Stavely, J.R., J.R. Steadman, D.P. Coyne, and D.T. Lindgren. 1989. BelNeb Rust Resistant-1 and -2 great northern dry bean germplasm. HortScience 24:400-401.

Varela, O., J.S. Beaver, M. Zapata, P. Miklas and S. Cianzaio. 1995. Evaluacion de frijol (*Phaseolus vulgaris* L.) en generaciones tempranas para resistencia a *Xanthomonas campestris* pv. *phaseoli* (Smith) Dye. J. of Agric. of the Univ. of Puerto Rico 80:55-63.
Yu, K., S. J. Park and V. Poysa. 2000. Marker-assisted selection of common beans for resistance to common bacterial blight: efficacy and economics. Plant Breedg. 119: 411-415.

Yu, K., S. J. Park, B. Zhang, M. Haffner and V. Poysa. 2000. An SSR marker in the nitrate reductase gene of common bean is tightly linked to a major gene controlling resistance to common bacterial blight. Euphytica 138: 89-95.

Zapata, M., G.F. Freytag and R.E. Wilkinson. 1985. Evaluation for bacterial blight resistance in beans. *Phytopathology* 75:1032-1039.

Zapata, M., G.F. Freytag and R.E. Wilkinson. 2004. Registration of five common bean germplasm lines resistant to common bacterial blight: W-BB-11, W-BB-20-1, W-BB-35, W-BB-52, and W-BB-11-56. *Annu. Rept. Bean Improv. Coop.* 47:333-337.